DROPLET & CONTACT PRECAUTIONS

Clean hands before entering and when leaving room

and

Please report to staff before entering

Bed #

ring and Ig room

STOP



Clean hands with A) hand foam/gel or B) soap and water

Required:

- Point of Care Risk Assessment
- Gown & Gloves
- Procedure mask with eye protection
 When within 2 metres of patient
- Keep 2 metres between patients





Families

visitors:

KEEP SIGN POSTED UNTIL ROOM CLEANED HOUSEKEEPER will remove sign after cleaning

9-90714





Point of Care Risk Assessment

Risk	Protection	
Contact with patient or environment expected	Hand hygiene	
 Splash or spray of blood or body fluids/ secretions anticipated 	Mask and eye protectionPut on gown if soiling of clothing is likely	
 Contact with mucous membranes Non-intact skin, blood, body fluids, secretions, excretions or soiled or likely soiled item/surface anticipated 	 Perform hand hygiene, then don gloves Perform hand hygiene after PPE removal and before leaving patient environment 	

Additional Precautions Guidelines

	Contact	Contact Plus	Droplet + Contact	Airborne	Airborne+ Contact	
Syndromic precautions	Draining wound, diarrhea, infestation	Diarrhea (<i>C.difficile</i> positive)	 Any of: Vomiting, with or without diarrhea New/worsening cough Fever, with or without rash Toxic shock 	Fever + weight loss + cough + high risk for TB	Disseminated rash + fever	
Private room	Preferred. For suspect & confirmed CPO: yes	Preferred	Preferred. If in multi- bed room, draw curtain.	Yes	Yes	
Negative pressure room	No	No	No	Yes	Yes	
Staff PPE	Gown + gloves	Gown + gloves	Procedure mask + eye protection + gown + gloves	N95 respirator	N95 respirator + gown + gloves	
Visitor PPE	Gown + gloves if direct care	Gown + gloves	Procedure mask + eye protection (+gown +gloves if direct care)	Offer N95 respirator to visitor	N95 respirator (+gown +gloves if direct care)	
Parents of pediatric patients	Clean hands before entering and on leaving room. Do not go into common areas such as patient kitchens, playrooms, school rooms, or patient lounges.					
Patient wears a procedure mask during transport	No	No	Yes	Yes	Yes	